

# On Partitions of Finite Vector Spaces of Low Dimension Over $GF(2)$

S.I. El-Zanati, G.F. Seelinger, P.A. Sissokho,  
L.E. Spence, and C. Vanden Eynden

4520 Mathematics Department  
Illinois State University  
Normal, Illinois 61790–4520, U.S.A.

## Abstract

Let  $V_n(q)$  denote a vector space of dimension  $n$  over the field with  $q$  elements. A set  $\mathcal{P}$  of subspaces of  $V_n(q)$  is a *partition* of  $V_n(q)$  if every nonzero vector in  $V_n(q)$  is contained in exactly one subspace of  $\mathcal{P}$ . If there exists a partition of  $V_n(q)$  containing  $a_i$  subspaces of dimension  $n_i$  for  $1 \leq i \leq k$ , then  $(a_k, a_{k-1}, \dots, a_1)$  must satisfy the Diophantine equation  $\sum_{i=1}^k a_i(q^{n_i} - 1) = q^n - 1$ . In general, however, not every solution of this Diophantine equation corresponds to a partition of  $V_n(q)$ . In this article, we determine all solutions of the Diophantine equation for which there is a corresponding partition of  $V_n(2)$  for  $n \leq 7$ .

## 1 Introduction

Let  $K = GF(q)$  be the finite field with  $q$  elements and  $V = V_n(q)$  be a vector space of dimension  $n$  over  $K$ . We say that a set  $\mathcal{P} = \{V_i\}_{i=1}^k$  of subspaces of  $V$  is a *partition* of  $V$  if and only if  $V = \cup_{i=1}^k V_i$  and  $V_i \cap V_j = \{0\}$  when  $i \neq j$ .

Suppose that  $\mathcal{P} = \{V_i\}_{i=1}^k$  is a partition of  $V$ , and let  $n_i = \dim V_i$ . It is easy to see that the following conditions are satisfied:

$$q^n - 1 = \sum_1^k (q^{n_i} - 1) \quad (1)$$

and

$$n_i + n_j \leq n \text{ when } i \neq j. \quad (2)$$

Condition (1) is true because every nonzero vector in  $V_n(q)$  belongs to exactly one of the subspaces  $V_i$ , and (2) holds because  $V_i \oplus V_j$  is a subspace of  $V$  with dimension  $n_i + n_j$ . Let  $W$  be a subspace of  $V_n(q)$  having dimension  $n - 1$ . Then  $\{V_i \cap W\}_{i=1}^k$  is a partition of  $W$ , which we call the *partition of  $W$  induced by  $\mathcal{P}$* , and, for  $i = 1, 2, \dots, k$ , we have  $\dim(V_i \cap W) = \dim V_i$  if  $V_i \subseteq W$ , and  $\dim(V_i \cap W) = \dim V_i - 1$  otherwise.

Bu [5] presents a number of sufficient conditions for the existence of certain partitions of  $V_n(q)$ . The first of these is a well-known result.

**Lemma 1.1.** *If  $m$  is a divisor of  $n$  and  $k = (q^n - 1)/(q^m - 1)$ , there exists a partition of  $V_n(q)$  consisting of  $k$  subspaces of dimension  $m$ .*

**Lemma 1.2.** For  $1 \leq d < \frac{1}{2}n$ , there exists a partition of  $V_n(q)$  consisting of 1 subspace of dimension  $n - d$  and  $q^{n-d}$  subspaces of dimension  $d$ .

**Lemma 1.3.** If  $n = ks - 1$ , where  $s > 1$ , there exists a partition of  $V_n(q)$  consisting of  $q^{(k-1)s}$  subspaces of dimension  $s - 1$  and  $(q^{(k-1)s} - 1)/(q^s - 1)$  subspaces of dimension  $s$ .

Other authors have considered variations on this problem. Let  $T = \{t_1, t_2, \dots, t_k\}$  be a set of positive integers with

$$2 \leq t_1 < t_2 < \dots < t_k = t. \quad (3)$$

A partition  $\mathcal{P}$  of  $V_n(q)$  is called a  $T$ -partition if (i) for every  $\alpha \in T$ , there is a  $U \in \mathcal{P}$  with  $\dim U = \alpha$ ; and (ii)  $\dim U \in T$  for any  $U \in \mathcal{P}$ . Beutelspacher proved in [1] that if  $t_1 = 2$ , then  $V_{2t}(q)$  has a  $T$ -partition. Heden shows in [6] that  $V_{2t}(q)$  has a  $T$ -partition for any arbitrary set of integers satisfying (3).

Before proceeding, we note that this partition problem also relates to the problem of finding optimal partial spreads and has applications to byte error control codes. For further information, we direct the interested reader to the article by Clark and Dunning [3], and to the references therein.

Suppose there exists a partition of  $V_n(q)$  into  $a_i$  subspaces of dimension  $i$  for  $1 \leq i \leq k$ . By (1),  $(a_1, a_2, \dots, a_k)$  must satisfy the Diophantine equation

$$\sum_{i=1}^k a_i (q^{n_i} - 1) = q^n - 1. \quad (4)$$

In general, however, not every solution of (4) corresponds to a partition of  $V_n(q)$ . For example, Corollary 2.2 below shows that there is no partition of  $V_5(2)$  into 10 subspaces of dimension 2 and 1 subspace of dimension 1. In this article, we classify all solutions of (4) for which there is a corresponding partition of  $V_n(2)$  when  $n \leq 7$ .

## 2 Feasible Partition Types

We begin by proving a third necessary condition for a solution to (4) to correspond to a partition of  $V_n(2)$ .

**Lemma 2.1.** Let  $V$  be a finite dimensional vector space over a finite field  $F$ . Then  $\sum_{v \in V} v = 0$  if  $\dim V \geq 2$  or  $\text{char } F \neq 2$ .

*Proof.* If  $\text{char } F \neq 2$ , then every nonzero vector  $v \in V$  has a unique additive inverse  $-v \neq v$ . So, in this case, it is clear that  $\sum_{v \in V} v = 0$ .

If  $\text{char } F = 2$ , then we can reduce to the case where  $F$  is the field with two elements. Let  $V$  be a finite-dimensional vector space over  $F$  of dimension  $n \geq 2$ , and let  $\{u_1, u_2, \dots, u_n\}$  be a basis for  $V$ . Define  $w = \sum_{v \in V} v$ . Then  $w = \sum_{i=1}^n \alpha_i u_i$  for some  $\alpha_i \in F$ .

We claim  $\alpha_i = 0$  for all  $i$ . Indeed, for any  $v \in V$ , we can write  $v = \sum_{i=1}^n \epsilon_{i,v} u_i$ . Therefore  $w = \sum_{v \in V} \sum_{i=1}^n \epsilon_{i,v} u_i = \sum_{i=1}^n (\sum_{v \in V} \epsilon_{i,v}) u_i$ , so that  $\alpha_i = \sum_{v \in V} \epsilon_{i,v}$ . In particular,  $\alpha_i = t(1_F)$ , where  $t$  is the number of vectors such that  $\epsilon_{i,v} = 1_F$ . But the set of vectors whose  $i$ th component is  $1_F$  is given by  $\{u_i + \sum_{j \neq i} \epsilon_j u_j : \epsilon_j \in F\}$ . Since the order of this set is even,  $t(1_F) = 0_F$ ; so  $\alpha_i = 0$  for all  $i$ . Therefore  $\sum_{v \in V} v = 0$ .  $\square$

**Corollary 2.2.** In any partition of  $V_n(q)$  ( $n \geq 2$ ) into subspaces, the sum of the vectors in the 1-dimensional subspaces of the partition is 0.

*Proof.* From Lemma 2.1, the sum of the vectors in  $V_n(q)$  is 0, and the same is true for the sum of the vectors in the subspaces of dimension greater than 1. The result follows.  $\square$

Note that Corollary 2.2 shows that there can be no partition of  $V_5(2)$  containing exactly ten subspaces of dimension 2 and one subspace of dimension 1. However, such a partition satisfies both of the necessary conditions (1) and (2). Thus Corollary 2.2 provides a new necessary condition for the existence of a partition of  $V_n(2)$ . Because every vector in  $V_n(2)$  is its own additive inverse, Corollary 2.2 implies that *if a partition of  $V_n(2)$  contains subspaces of dimension 1, then there cannot be fewer than 3 subspaces of dimension 1.*

We call a sequence of nonnegative integers  $\langle a_k, a_{k-1}, \dots, a_1 \rangle$  a *partition type* for  $V_n(2)$  and say that it is *feasible* if the following four conditions hold:

- (a)  $\sum_{i=1}^k a_i(2^i - 1) = 2^n - 1$ .
- (b) If both  $a_i$  and  $a_j$  are nonzero for  $i \neq j$ , then  $i + j \leq n$ .
- (c) If  $i > n/2$ , then  $a_i \leq 1$ .
- (d)  $a_1 \neq 1$  and  $a_1 \neq 2$ .

Thus a necessary condition for the existence of a partition of  $V_n(2)$  containing exactly  $a_i$  subspaces of dimension  $i$  for  $i = 1, 2, \dots, k$  is that the partition type  $\langle a_k, a_{k-1}, \dots, a_1 \rangle$  be feasible.

If there is a partition of  $V_n(2)$  containing exactly  $a_i$  subspaces of dimension  $i$  for  $i = 1, 2, \dots, k$ , then we say that the partition type  $\langle a_k, a_{k-1}, \dots, a_1 \rangle$  is *realized*. In Sections 3, 5, and 6 below, we determine, for  $n \leq 7$ , all of the feasible partition types for  $V_n(2)$  that are realized. Specifically, we prove the following result.

**Main Theorem.** *For  $n \leq 7$ , all the feasible partition types of  $V_n(2)$  are realized except  $\langle 7, 3, 5 \rangle$  when  $n = 6$  and  $\langle 1, 13, 7, 0 \rangle$ ,  $\langle 1, 13, 6, 3 \rangle$ ,  $\langle 1, 14, 3, 5 \rangle$ , and  $\langle 17, 1, 5 \rangle$  when  $n = 7$ .*

In a forthcoming article [4], we use this result to provide the base of a proof by mathematical induction showing that, for every integer  $n > 2$ , a feasible partition type of  $V_n(2)$  with the form  $\langle a_3, a_2, 0 \rangle$  is realized if and only if  $a_2 \neq 1$ . In another article [2], we show how partitions of  $V_n(q)$  relate to uniformly resolvable designs.

**Convention.** Throughout the remainder of this paper, if  $S$  is a subset of  $V_n(2)$  such that  $S \cup \{0\}$  is a subspace, we will use the term *subspace* to refer to  $S$ .

### 3 Partitions of $V_n(2)$ for $n \leq 5$

#### Partitions of $V_3(2)$

The 3 feasible partition types are  $\langle 1, 0, 0 \rangle$ ,  $\langle 1, 4 \rangle$ , and  $\langle 7 \rangle$ . Clearly all of these can be realized.

#### Partitions of $V_4(2)$

The 8 feasible partition types are  $\langle 1, 0, 0, 0 \rangle$ ,  $\langle 1, 0, 8 \rangle$ , and  $\langle 5 - i, 3i \rangle$  for  $0 \leq i \leq 5$ . Partitions of type  $\langle 1, 0, 0, 0 \rangle$  and  $\langle 5, 0 \rangle$  are realized by Lemma 1.1. A partition of type  $\langle 1, 0, 8 \rangle$  can be obtained by extracting a 3-dimensional subspace of  $V_4(2)$  and regarding the remaining vectors as the nonzero vectors in subspaces of dimension 1. The other partitions are obtained by regarding the nonzero vectors in 2-dimensional subspaces in a partition of type  $\langle 5, 0 \rangle$  as the nonzero vectors in 3 subspaces of dimension 1.

## Partitions of $V_5(2)$

The 21 feasible partition types are  $\langle 1, 0, 0, 0, 0 \rangle$ ,  $\langle 1, 0, 0, 16 \rangle$ , and  $\langle 1 - i, 8 + i - j, 4i + 3j \rangle$  for  $0 \leq j \leq 8 + i$  and  $0 \leq i \leq 1$ . Partitions of types  $\langle 1, 0, 0, 0, 0 \rangle$  and  $\langle 1, 0, 0, 16 \rangle$  clearly exist, and a partition of type  $\langle 1, 8, 0 \rangle$  can be realized by Lemma 1.2. The remaining partitions can be obtained from one of type  $\langle 1, 8, 0 \rangle$  by regarding the subspace of dimension 3 as 1 subspace of dimension 2 and 4 subspaces of dimension 1, or regarding some subspaces of dimension 2 as 3 subspaces of dimension 1.

## 4 Reconfiguring Subspaces

In Section 3, subspaces in a partition were changed into new subspaces to create a new partition. We call this process *reconfiguring* the subspaces. Reconfiguring subspaces plays an essential role in constructing partitions of  $V_6(2)$  and  $V_7(2)$ .

We have already seen that a subspace of dimension  $m$  can be reconfigured into a subspace of dimension  $k < m$  and subspaces of dimension 1. In addition, any subspace of dimension 4 can be reconfigured into 5 subspaces of dimension 2.

If  $n = 2k$ , then any three subspaces of dimension  $k$  in a partition of  $V_n(2)$  can be reconfigured into  $r = 2^k - 1$  subspaces of dimension 2. For suppose that  $V_1, V_2$ , and  $V_3$  are any three  $k$ -dimensional subspaces in a partition of  $V_n(2)$ . Let the nonzero elements in these subspaces be

$$x_1, x_2, \dots, x_r, \quad y_1, y_2, \dots, y_r, \quad \text{and} \quad z_1, z_2, \dots, z_r,$$

respectively. Since  $V_n(2) = V_1 \oplus V_2$ , it is possible to number the elements of  $V_1, V_2$ , and  $V_3$  so that  $x_i + y_i = z_i$  for  $i = 1, 2, \dots, r$ . Thus  $V_1, V_2$ , and  $V_3$  can be reconfigured into the 2-dimensional subspaces

$$\{0, x_1, y_1, z_1\}, \quad \{0, x_2, y_2, z_2\}, \quad \dots, \quad \{0, x_r, y_r, z_r\}.$$

In particular, any 3 subspaces of dimension 3 in a partition of  $V_6(2)$  can be reconfigured into 7 subspaces of dimension 2.

## 5 Partitions of $V_6(2)$

Clearly, only one feasible partition type for  $V_6(2)$  contains a subspace of dimension 6, and, because of (b) in the definition, only one feasible partition type for  $V_6(2)$  contains a subspace of dimension 5. A feasible partition type that contains a subspace of dimension 4 must be a nonnegative integer solution of the system

$$\begin{aligned} 15a_4 + 7a_3 + 3a_2 + a_1 &= 63 \\ a_4 &= 1 \\ a_3 &= 0, \end{aligned}$$

where  $a_1 \neq 1$  and  $a_1 \neq 2$ , that is, a nonnegative integer solution of the equation  $3a_2 + a_1 = 48$  such that  $a_1 \neq 1$  and  $a_1 \neq 2$ . We can regard these feasible partition types as solutions of the inequality  $3a_2 \leq 48$ , and so there are 17 feasible partition types containing a subspace of dimension 4.

As above, we can regard a feasible partition type having no subspaces of dimension greater than 3 as a nonnegative integer solution of  $7a_3 + 3a_2 \leq 63$  such that  $63 - (7a_3 + 3a_2) \neq 1$  and  $63 - (7a_3 + 3a_2) \neq 2$ . A straightforward count shows that there are 106 such solutions, and hence

Category	Type	
1	$\langle 1, 0, 0, 0, 0, 0 \rangle$	
2	$\langle 1, 0, 0, 0, 32 \rangle$	
3	$\langle 1, 0, 16 - j, 3j \rangle$	$0 \leq j \leq 16$
4	$\langle 9 - 3i, 7i - j, 3j \rangle$	$0 \leq j \leq 7i$ and $0 \leq i \leq 3$
5	$\langle 8 - 3i, 1 + 7i - j, 4 + 3j \rangle$	$0 \leq j \leq 1 + 7i$ and $0 \leq i \leq 2$
6	$\langle 7 - 3i, 3 + 7i - j, 5 + 3j \rangle$	$0 \leq j \leq 3 + 7i$ and $0 \leq i \leq 2$

Table 1: Feasible Partition Types of  $V_6(2)$

106 feasible partition types having no subspaces of dimension greater than 3. Thus, in all, there are 125 feasible partition types for  $V_6(2)$ , which are shown in Table 1.

Each partition in categories 3, 4, 5, and 6 can be obtained by reconfiguration from the one in the same category with  $i = j = 0$ . For example, a partition of type  $\langle 1, 0, 14, 6 \rangle$  (category 3) can be obtained from one of type  $\langle 1, 0, 16, 0 \rangle$  by reconfiguring 2 subspaces of dimension 2 into 6 subspaces of dimension 1. Also, a partition of type  $\langle 5, 4, 16 \rangle$  (category 5) can be obtained from one of type  $\langle 8, 1, 4 \rangle$  by reconfiguring 3 subspaces of dimension 3 into 7 subspaces of dimension 2 and then reconfiguring 4 subspaces of dimension 2 into 12 subspaces of dimension 1. In fact, Table 1 is arranged so that if a partition in some category is obtained, then all other partition types in that category having larger values of  $i$  or  $j$  can be realized in this manner. Hence, to verify which feasible partition types of  $V_6(2)$  can be realized, it is sufficient to show that one partition in each category exists and to prove that partition types in that category having smaller values of  $i$  or  $j$  cannot be realized.

Partitions in categories 1 and 2 clearly exist. A partition of type  $\langle 1, 0, 16, 0 \rangle$  (category 3) exists by Lemma 1.2. A partition of type  $\langle 9, 0, 0 \rangle$  (category 4) can be obtained from Lemma 1.1. A partition of type  $\langle 8, 1, 4 \rangle$  (category 5) can be obtained from a partition of type  $\langle 9, 0, 0 \rangle$  by reconfiguring 1 subspace of dimension 3 into 1 subspace of dimension 2 and 4 subspaces of dimension 1. Finally, Proposition 5.1 below shows that a partition of type  $\langle 7, 3, 5 \rangle$  (category 6) does not exist. However, a partition of type  $\langle 7, 2, 8 \rangle$  can be obtained from a partition of type  $\langle 9, 0, 0 \rangle$  by reconfiguring 2 subspaces of dimension 3 into 2 subspaces of dimension 2 and 8 subspaces of dimension 1, and a partition of type  $\langle 4, 10, 5 \rangle$  is constructed in Example 2 below.

**Proposition 5.1.** *There is no partition of  $V_6(2)$  of type  $\langle 7, 3, 5 \rangle$ .*

*Proof.* Suppose that  $\mathcal{P}$  is a partition of  $V_6(2)$  of type  $\langle 7, 3, 5 \rangle$ . Select a 5-dimensional subspace  $W$  of  $V_6(2)$  that contains all of the 1-dimensional subspaces in  $\mathcal{P}$ . By condition (c), the partition of  $W$  induced by  $\mathcal{P}$  is of type  $\langle a_3, a_2, a_1 \rangle$ , where

(i)  $a_3 = 1$ ,  $a_2 + a_1 = 14$ , and  $3a_2 + 1a_1 = 24$ , or

(ii)  $a_3 = 0$ ,  $a_2 + a_1 = 15$ , and  $3a_2 + 1a_1 = 31$ .

In case (i), we have  $a_2 = 5$  and  $a_1 = 9$ , which is a contradiction because there can be at most  $3 + 5 = 8$  subspaces of dimension 1 in the induced partition of  $W$ . In case (ii), we have  $a_2 = 8$

and  $a_1 = 7$ . For this to occur, there must be two nonzero vectors  $u$  and  $v$  that are contained in distinct 2-dimensional subspaces of  $\mathcal{P}$  and are contained in 1-dimensional subspaces of the induced partition of  $W$ . But the sum of the 5 nonzero vectors in the 1-dimensional subspaces of  $\mathcal{P}$  and the sum of the 7 nonzero vectors in the 1-dimensional subspaces of the induced partition of  $W$  must be 0. So  $u + v = 0$ , a contradiction. Thus there can be no partition of  $V_6(2)$  of type  $\langle 7, 3, 5 \rangle$ .  $\square$

Although a partition of type  $\langle 9, 0, 0 \rangle$  exists by Lemma 1.1, we now describe an explicit construction of such a partition that is used in the rest of this paper.

**Example 1** In what follows, we always identify a subspace by the set of its nonzero elements. Let  $V$  be the set of nonzero elements of a 3-dimensional subspace of  $V_n(2)$  with basis  $\{v_1, v_2, v_3\}$ . Then

$$V = \{v_1, v_2, v_3, v_4 = v_1 + v_2, v_5 = v_2 + v_3, v_6 = v_1 + v_2 + v_3, v_7 = v_1 + v_3\}.$$

For any integer  $1 \leq k \leq 7$ , define  $V^k$  to be the 7-tuple generated by the elements of  $V$  with first entry  $v_k$ , that is,

$$V^k = (v_k, v_{k+1}, \dots, v_{k+6}),$$

where the addition of indices is done modulo 7. If we do not need to specify the first entry of this  $k$ -tuple, then we simply write  $[V]$ .

Consider the recurrence relation

$$v_{i+3} = v_i + v_{i+1},$$

with the addition of indices performed modulo 7.

The sequence defined by this recurrence relation yields  $[V]$ . In particular, we have

$$\begin{aligned} v_4 &= v_1 + v_2, & v_5 &= v_2 + v_3, & v_6 &= v_3 + v_4 = v_1 + v_2 + v_3, \\ v_7 &= v_4 + v_5 = v_1 + v_3, & v_1 &= v_5 + v_6, & v_2 &= v_6 + v_7, & v_3 &= v_7 + v_8 \dots \end{aligned}$$

An addition table for  $V$  is shown in Table 2.

+	<b>0</b>	<b>v<sub>1</sub></b>	<b>v<sub>2</sub></b>	<b>v<sub>3</sub></b>	<b>v<sub>4</sub></b>	<b>v<sub>5</sub></b>	<b>v<sub>6</sub></b>	<b>v<sub>7</sub></b>
<b>0</b>	0	$v_1$	$v_2$	$v_3$	$v_4$	$v_5$	$v_6$	$v_7$
<b>v<sub>1</sub></b>	$v_1$	0	$v_4$	$v_7$	$v_2$	$v_6$	$v_5$	$v_3$
<b>v<sub>2</sub></b>	$v_2$	$v_4$	0	$v_5$	$v_1$	$v_3$	$v_7$	$v_6$
<b>v<sub>3</sub></b>	$v_3$	$v_7$	$v_5$	0	$v_6$	$v_2$	$v_4$	$v_1$
<b>v<sub>4</sub></b>	$v_4$	$v_2$	$v_1$	$v_6$	0	$v_7$	$v_3$	$v_5$
<b>v<sub>5</sub></b>	$v_5$	$v_6$	$v_3$	$v_2$	$v_7$	0	$v_1$	$v_4$
<b>v<sub>6</sub></b>	$v_6$	$v_5$	$v_7$	$v_4$	$v_3$	$v_1$	0	$v_2$
<b>v<sub>7</sub></b>	$v_7$	$v_3$	$v_6$	$v_1$	$v_5$	$v_4$	$v_2$	0

Table 2: An Addition Table for  $V$

Now let  $U$  and  $W$  be subspaces of  $V_n(2)$  having dimension 3 such that  $U \cap W = \{0\}$ . As above, the nonzero elements  $u_1, u_2, \dots, u_7$  and  $w_1, w_2, \dots, w_7$  of  $U$  and  $W$ , respectively, can be chosen to satisfy the recurrence relations

$$u_{i+3} = u_i + u_{i+1} \quad \text{and} \quad w_{i+3} = w_i + w_{i+1},$$

where addition of indices is done modulo 7. The elements of  $U$  and  $W$  can be added as in Table 2.

Define

$$A_{i-1} = \{u_1 + w_i, u_2 + w_{i+1}, \dots, u_7 + w_{i+6}\},$$

for  $1 \leq i \leq 7$ . Then  $A_{i-1}$  is the set of nonzero elements from a subspace of dimension 3 with basis

$$\{u_1 + w_i, u_2 + w_{i+1}, u_3 + w_{i+2}\}.$$

For  $1 \leq k \leq 7$ , let

$$U^k = (u_k, u_{k+1}, \dots, u_{k+6}) \quad \text{and} \quad W^k = (w_k, w_{k+1}, \dots, w_{k+6}),$$

and for  $0 \leq i \leq 6$ , let

$$A_i^k = (u_k + w_{i+k}, u_{k+1} + w_{i+k+1}, \dots, u_{k+6} + w_{i+k+6}).$$

The subspaces  $U$ ,  $W$ , and  $A_i \cup \{0\}$  ( $0 \leq i \leq 6$ ) form a partition of

$$U \oplus W = \{u + w : u \in U \text{ and } w \in W\}$$

into 9 subspaces of dimension 3. Note that the 7-tuples  $U^k$  and  $W^k$  satisfy the same recurrence relation as the vectors  $u_k$  and  $w_k$ , that is,

$$U^k + U^{k+1} = U^{k+3} \quad \text{and} \quad W^k + W^{k+1} = W^{k+3}.$$

Moreover, we have  $A_i^k = U^k + W^{i+k}$ .

□

### Example 2

We prove the existence of a partition of  $V_6(2)$  of type  $\langle 4, 10, 5 \rangle$  by showing that the 3-dimensional subspaces  $A_0, A_1, A_2, A_3$ , and  $A_4$  in Example 1 can be reconfigured into 10 subspaces of dimension 2 and 5 subspaces of dimension 1. The nonzero vectors in the subspaces of dimension 2 are:

$$\begin{aligned} &\{u_1 + w_1, u_2 + w_3, u_4 + w_7\}, && \{u_2 + w_2, u_6 + w_1, u_7 + w_4\}, \\ &\{u_3 + w_3, u_4 + w_5, u_6 + w_2\}, && \{u_4 + w_4, u_6 + w_7, u_3 + w_5\}, \\ &\{u_5 + w_5, u_7 + w_1, u_4 + w_6\}, && \{u_6 + w_6, u_2 + w_4, u_7 + w_3\}, \\ &\{u_7 + w_7, u_1 + w_2, u_3 + w_6\}, && \{u_3 + w_4, u_5 + w_7, u_2 + w_5\}, \\ &\{u_5 + w_6, u_1 + w_4, u_6 + w_3\}, & \text{and} & \{u_1 + w_5, u_2 + w_6, u_4 + w_1\}. \end{aligned}$$

Together with the 3-dimensional subspaces  $W$ ,  $U$ ,  $A_5$ , and  $A_6$ , these 10 subspaces of dimension 2 and the remaining 5 subspaces of dimension 1 form a partition of  $V_6(2)$  of type  $\langle 4, 10, 5 \rangle$ . □

The results obtained in this section show that every feasible partition type for  $V_6(2)$  can be realized except  $\langle 7, 3, 5 \rangle$ .

## 6 Partitions of $V_7(2)$

### 6.1 Feasible Partition Types of $V_7(2)$

By an analysis similar to that in Section 5, it can be shown that there are 758 feasible partition types for  $V_7(2)$ , which are shown in Table 3.

Category	Partition Type		Example	Category	Partition Type		Example
1	$\langle 1, 0, 0, 0, 0, 0, 0 \rangle$						
2	$\langle 1, 0, 0, 0, 0, 64 \rangle$						
3	$\langle 1, 0, 0, 32 - j, 3j \rangle$	$0 \leq j \leq 32$		21	$\langle 17, 1 - j, 5 + 3j \rangle$	$0 \leq j \leq 1$	
4	$\langle 1, 16, 0, 0 \rangle$			22	$\langle 16, 5 - j, 3j \rangle$	$0 \leq j \leq 5$	
5	$\langle 1, 15, 1 - j, 4 + 3j \rangle$	$0 \leq j \leq 1$	5	23	$\langle 15, 6 - j, 4 + 3j \rangle$	$0 \leq j \leq 6$	
6	$\langle 1, 14, 3 - j, 5 + 3j \rangle$	$0 \leq j \leq 3$	5	24	$\langle 14, 8 - j, 5 + 3j \rangle$	$0 \leq j \leq 8$	8
7	$\langle 1, 13, 7 - j, 3j \rangle$	$0 \leq j \leq 7$	4	25	$\langle 13, 12 - j, 3j \rangle$	$0 \leq j \leq 12$	9
8	$\langle 1, 12, 8 - j, 4 + 3j \rangle$	$0 \leq j \leq 8$	5	26	$\langle 12, 13 - j, 4 + 3j \rangle$	$0 \leq j \leq 13$	
9	$\langle 1, 11, 10 - j, 5 + 3j \rangle$	$0 \leq j \leq 10$	6	27	$\langle 11, 15 - j, 5 + 3j \rangle$	$0 \leq j \leq 15$	
10	$\langle 1, 10, 14 - j, 3j \rangle$	$0 \leq j \leq 14$	3	28	$\langle 10, 19 - j, 3j \rangle$	$0 \leq j \leq 19$	
11	$\langle 1, 9, 15 - j, 4 + 3j \rangle$	$0 \leq j \leq 15$	5	29	$\langle 9, 20 - j, 4 + 3j \rangle$	$0 \leq j \leq 20$	
12	$\langle 1, 8, 17 - j, 5 + 3j \rangle$	$0 \leq j \leq 17$	7	30	$\langle 8, 22 - j, 5 + 3j \rangle$	$0 \leq j \leq 22$	
13	$\langle 1, 7, 21 - j, 3j \rangle$	$0 \leq j \leq 21$	3	31	$\langle 7, 26 - j, 3j \rangle$	$0 \leq j \leq 26$	
14	$\langle 1, 6, 22 - j, 4 + 3j \rangle$	$0 \leq j \leq 22$	5	32	$\langle 6, 27 - j, 4 + 3j \rangle$	$0 \leq j \leq 27$	
15	$\langle 1, 5, 24 - j, 5 + 3j \rangle$	$0 \leq j \leq 24$	6	33	$\langle 5, 29 - j, 5 + 3j \rangle$	$0 \leq j \leq 29$	
16	$\langle 1, 4, 28 - j, 3j \rangle$	$0 \leq j \leq 28$	3	34	$\langle 4, 33 - j, 3j \rangle$	$0 \leq j \leq 33$	
17	$\langle 1, 3, 29 - j, 4 + 3j \rangle$	$0 \leq j \leq 29$	5	35	$\langle 3, 34 - j, 4 + 3j \rangle$	$0 \leq j \leq 34$	
18	$\langle 1, 2, 31 - j, 5 + 3j \rangle$	$0 \leq j \leq 31$	6	36	$\langle 2, 36 - j, 5 + 3j \rangle$	$0 \leq j \leq 36$	
19	$\langle 1, 1, 35 - j, 3j \rangle$	$0 \leq j \leq 35$	3	37	$\langle 1, 40 - j, 3j \rangle$	$0 \leq j \leq 40$	
20	$\langle 1, 0, 36 - j, 4 + 3j \rangle$	$0 \leq j \leq 36$	5	38	$\langle 41 - j, 4 + 3j \rangle$	$0 \leq j \leq 41$	

Table 3: Feasible Partition Types of  $V_7(2)$

We will show that all of the partition types in Table 3 can be realized except for four:  $\langle 1, 14, 3, 5 \rangle$  (category 6),  $\langle 1, 13, 7, 0 \rangle$  (category 7),  $\langle 1, 13, 6, 3 \rangle$  (category 7), and  $\langle 17, 1, 5 \rangle$  (category 21).

Note that Table 3 is arranged so that if a partition in some category is obtained, then all other partition types in that category having larger values of  $j$  can be realized by reconfiguring 2-dimensional subspaces into subspaces of dimension 1. For example, the partition in category 3 of type  $\langle 1, 0, 0, 32 - j, 3j \rangle$  can be obtained from one of type  $\langle 1, 0, 0, 32, 0 \rangle$  by regarding  $j$  subspaces of dimension 2 as  $3j$  subspaces of dimension 1.

### 6.2 Partition Types of $V_7(2)$ That Cannot Be Realized

In this subsection, we prove that 4 of the feasible partition types of  $V_7(2)$  cannot be realized.



**Lemma 6.1.** *Let  $\mathcal{B}$  be a basis for an  $n$ -dimensional vector space  $V$ , and let  $\mathcal{B}'$  be a proper subset of  $\mathcal{B}$ . Then  $V$  has an  $(n - 1)$ -dimensional subspace  $W$  such that  $W \cap \mathcal{B} = \mathcal{B}'$ .*

*Proof.* We can assume that  $\mathcal{B}$  is the basis  $\{e_1, e_2, \dots, e_n\}$  and  $\mathcal{B}' = \{e_1, e_2, \dots, e_k\}$  for some  $k < n$ , where  $e_i$  is the vector whose  $i$ th component is 1 and whose other components are 0. The desired subspace is  $W = \{(\alpha_1, \alpha_2, \dots, \alpha_n) \in V : \alpha_{k+1} + \alpha_{k+2} + \dots + \alpha_n = 0\}$ .  $\square$

**Proposition 6.2.** *No partition of  $V_7(2)$  of type  $\langle 1, 13, 7, 0 \rangle$  exists.*

*Proof.* Assume that  $\mathcal{P}$  is a partition of type  $\langle 1, 13, 7, 0 \rangle$ . By (2), the partition of a 6-dimensional subspace of  $V_7(2)$  induced by  $\mathcal{P}$  is of type  $\langle 1, 0, a_2, a_1 \rangle$ , where  $a_2$  and  $a_1$  satisfy

$$\begin{aligned} a_2 + a_1 &= 20 \\ 3a_2 + a_1 &= 48, \end{aligned}$$

or of type  $\langle a_3, a_2, a_1 \rangle$ , where  $a_3$ ,  $a_2$ , and  $a_1$  satisfy

$$\begin{aligned} a_3 + a_2 + a_1 &= 21 \\ 7a_3 + 3a_2 + a_1 &= 63. \end{aligned}$$

It is easily checked that there is no integral solution of either system in which  $a_1 = 7$ . Moreover, only the second system has an integral solution in which  $a_1 = 0$ , and it is  $a_3 = 0, a_2 = 21, a_1 = 0$ . However, there can be no induced partition of type  $\langle 21, 0 \rangle$  because there must be at least one subspace of dimension 3 in the induced partition. Thus no partition induced by  $\mathcal{P}$  can contain 0 or 7 subspaces of dimension 1.

Let  $V_1, V_2, \dots, V_7$  denote the 2-dimensional subspaces in  $\mathcal{P}$ , and let  $S = \{u_1, u_2, \dots, u_k\}$  be a linearly independent subset of  $V_1 \cup V_2 \cup \dots \cup V_7$  that is maximal with respect to the property that each  $u_i$  is in a different 2-dimensional subspace in  $\mathcal{P}$ . By renumbering, if necessary, we may assume that  $u_i \in V_i$  for  $i = 1, 2, \dots, k$ .

*Case 1:  $k = 7$ .*

In this case,  $S$  is a basis for  $V_7(2)$ . Apply Lemma 6.1 to find a 6-dimensional subspace  $W$  of  $V_7(2)$  that contains none of the vectors in  $S$ . Then  $W$  does not contain any  $V_i$ , and hence the partition of  $W$  induced by  $\mathcal{P}$  must contain 7 subspaces of dimension 1, a contradiction.

*Case 2:  $k < 7$ .*

We claim that  $V_i \subseteq \text{Span } S$  for  $k < i \leq 7$ . For otherwise, there is a vector  $u_i$  in  $V_i$  that is not in  $\text{Span } S$ . But then the set  $\{u_1, u_2, \dots, u_k, u_i\}$  contradicts the maximality of  $S$ .

Let  $W$  be a 6-dimensional subspace of  $V_7(2)$  containing  $S$ . Since  $V_i \subseteq \text{Span } S$  for  $k < i \leq 7$ , the 1-dimensional subspaces in the partition of  $W$  induced by  $\mathcal{P}$  must be  $V_i \cap W = \{0, u_i\}$ , where  $1 \leq i \leq k$ . But the sum of the vectors in the 1-dimensional subspaces of this partition must be 0; so we have a contradiction of the linear independence of  $S$ .

Since each case leads to a contradiction, we conclude that there is no partition of type  $\langle 1, 13, 7, 0 \rangle$ .  $\square$

**Corollary 6.3.** *No partition of  $V_7(2)$  of type  $\langle 1, 13, 6, 3 \rangle$  exists.*

*Proof.* Suppose that  $\mathcal{P}$  is a partition of  $V_7(2)$  of type  $\langle 1, 13, 6, 3 \rangle$ . Because the vectors in the subspaces of  $\mathcal{P}$  of dimension 1 must sum to 0, the union of these subspaces must be a subspace of dimension 2. This yields a partition of type  $\langle 1, 13, 7, 0 \rangle$ , in contradiction of Proposition 6.2.  $\square$

**Proposition 6.4.** *No partition of  $V_7(2)$  of type  $\langle 1, 14, 3, 5 \rangle$  exists.*

*Proof.* Let  $\mathcal{P}$  be a partition of  $V_7(2)$  of type  $\langle 1, 14, 3, 5 \rangle$ , and let  $u_1, u_2, \dots, u_5$  denote the nonzero vectors in the five subspaces of dimension 1. If these were all contained in a subspace  $S$  of dimension 3, then, by Lemma 2.1 and Corollary 2.2, both the sum of the vectors in  $S$  and the sum of  $u_1, u_2, \dots, u_5$  would be 0. But then the sum of the other two nonzero vectors in  $S$  would be 0, which is impossible. Thus we can assume that  $u_1, u_2, u_3$ , and  $u_4$  are linearly independent.

Let  $W$  be a 6-dimensional subspace of  $V_7(2)$  that contains each of  $u_1, u_2, \dots, u_5$ . The partition of  $W$  induced by  $\mathcal{P}$  is of type  $\langle 1, 0, a_2, a_1 \rangle$ , where

$$\begin{aligned} a_2 + a_1 &= 22 \\ 3a_2 + a_1 &= 48 \end{aligned}$$

or of type  $\langle a_3, a_2, a_1 \rangle$ , where

$$\begin{aligned} a_3 + a_2 + a_1 &= 23 \\ 7a_3 + 3a_2 + a_1 &= 63. \end{aligned}$$

The unique solution of the first system is  $a_2 = 13$ ,  $a_1 = 9$ . However, the partition of  $W$  induced by  $\mathcal{P}$  cannot contain more than  $3 + 5 = 8$  subspaces of dimension 1, a contradiction.

The integral solutions of the second system with  $5 \leq a_1 \leq 8$  are  $a_3 = 1$ ,  $a_2 = 17$ ,  $a_1 = 5$  and  $a_3 = 2$ ,  $a_2 = 14$ ,  $a_1 = 7$ . In the latter case, the sum of the vectors in the 5 subspaces of dimension 1 in  $\mathcal{P}$  and the sum of the vectors in the 7 subspaces of dimension 1 in the partition of  $W$  induced by  $\mathcal{P}$  are both 0. Thus the 2 nonzero vectors in the 2 additional 1-dimensional subspaces of the partition of  $W$  must sum to 0, which is impossible. Therefore the partition of  $W$  induced by  $\mathcal{P}$  must be of type  $\langle 1, 17, 5 \rangle$ .

Let  $S' = \text{Span} \{u_1, u_2, u_3, u_4\}$ . If each of the 2-dimensional subspaces of  $\mathcal{P}$  were contained in  $S'$ , the 9 nonzero vectors in these subspaces and the vectors  $u_1, u_2, \dots, u_5$  would account for 14 of the 15 nonzero vectors in  $S'$ . Moreover, the sum of these 14 vectors is 0, and so the sum of the vectors in  $S'$  cannot be 0, which is impossible. Hence some element  $v$  in a 2-dimensional subspace  $V'$  of  $\mathcal{P}$  is not in  $S$ , and so the set  $\{u_1, u_2, u_3, u_4, v\}$  is linearly independent. By Lemma 6.1, there is a 6-dimensional subspace  $W'$  of  $V_7(2)$  that contains  $u_1, u_2, u_3$ , and  $u_4$ , but not  $v$ . Because  $u_5 = u_1 + u_2 + u_3 + u_4$ ,  $W'$  also contains  $u_5$ . Thus  $W'$  contains each of the 1-dimensional subspaces of  $\mathcal{P}$  but not the 2-dimensional subspace  $V'$ , and so the partition of  $W'$  induced by  $\mathcal{P}$  contains at least 6 subspaces of dimension 1, a contradiction. It follows that there is no partition of  $V_7(2)$  of type  $\langle 1, 14, 3, 5 \rangle$ .  $\square$

**Proposition 6.5.** *No partition of  $V_7(2)$  of type  $\langle 17, 1, 5 \rangle$  exists.*

*Proof.* Assume that  $\mathcal{P}$  is a partition of  $V_7(2)$  of type  $\langle 17, 1, 5 \rangle$ . As in the proof of Proposition 6.4, there are 4 linearly independent vectors  $u_1, u_2, u_3, u_4$  in the 1-dimensional subspaces. Extend  $\{u_1, u_2, u_3, u_4\}$  to a basis for  $V_7(2)$ , and apply Lemma 6.1 to obtain a 6-dimensional subspace  $W$  of  $V_7(2)$  that contains none of  $u_1, u_2, u_3$ , and  $u_4$ . The complement of  $W$  in  $V_7(2)$  can contain no 2-dimensional subspace, and because  $u_1$  and  $u_2$  are not in  $W$ , it follows that  $u_1 + u_2$  is in  $W$ . Similarly,  $u_3 + u_4$  is in  $W$ , and so  $u_5 = (u_1 + u_2) + (u_3 + u_4)$  is in  $W$ . Therefore the partition of  $W$  induced by  $\mathcal{P}$  contains either one or two subspaces of dimension 1, a contradiction of Corollary 2.2. Hence there is no partition of type  $\langle 17, 1, 5 \rangle$ .  $\square$

### 6.3 Partitions of $V_7(2)$ Containing a Subspace of Dimension 4 or More

Partition types of  $V_7(2)$  that contain a subspace of dimension 5 or more are listed in categories 1, 2, and 3 of Table 3. Because a partition of type  $\langle 1, 0, 0, 32, 0 \rangle$  exists by Lemma 1.2, all the partition types in categories 1, 2, and 3 can be realized.

Although a partition of type  $\langle 1, 16, 0, 0 \rangle$  (category 4) is known to exist by Lemma 1.2, we now construct a particular partition of this type that is useful in our examples. This construction uses the notation of Example 1 and the addition in Table 2, where  $U$  and  $W$  denote 3-dimensional subspaces of  $V_7(2)$  such that  $U \cap W = \{0\}$ . Choose an element  $x$  of  $V_7(2)$  not in  $U \oplus W$ , and let  $W^*$  denote the set of nonzero elements in  $W$ . Then the sets  $\{0, x\} \oplus U$ ,  $W^*$ ,  $x + W^*$ ,  $A_i$ , and  $x + A_i$  ( $0 \leq i \leq 6$ ) form a partition of the set  $V_7(2)$  into 17 subsets in which each set other than  $\{0, x\} \oplus U$  contains precisely 7 elements.

For any 7-tuple  $S = (s_1, s_2, \dots, s_7)$ , denote by  $x + S$  the 7-tuple

$$x + S = (x + s_1, x + s_2, \dots, x + s_7).$$

By using the ideas in Example 1 and Table 2, we see that the following equations are true.

$$(x + W^1) + (x + A_0^6) + A_1^1 + A_6^6 + (x + A_3^1) + (x + A_2^5) = A_5^5 \quad (5)$$

$$W^1 + A_0^2 + (x + A_1^6) + A_2^2 + (x + A_4^6) + (x + A_6^7) = (x + A_7^7) \quad (6)$$

We can use equations (5) and (6) to create 16 subspaces of dimension 3 from the 7-tuples  $[W]$ ,  $x + [W]$ ,  $[A_i]$ , and  $x + [A_i]$  ( $0 \leq i \leq 6$ ). In Table 4, the respective components of the 7-tuples in equation (5) are listed vertically. Using the recurrence relations defined in Example 1, we see that each set  $S_i$  consisting of the  $i$ th components of these 7-tuples is a 3-dimensional subspace of  $V_7(2)$ . Thus equation (5) yields 7 subspaces of  $V_7(2)$  of dimension 3. Similarly, equation (6) yields 7 subspaces of  $V_7(2)$  of dimension 3.

	$x + W^1$	$x + A_0^6$	$A_1^1$	$A_6^6$	$x + A_3^1$	$x + A_2^5$	$A_5^5$
$S_1$	$x + w_1$	$x + u_6 + w_6$	$u_1 + w_2$	$u_6 + w_5$	$x + u_1 + w_4$	$x + u_5 + w_7$	$u_5 + w_3$
$S_2$	$x + w_2$	$x + u_7 + w_7$	$u_2 + w_3$	$u_7 + w_6$	$x + u_2 + w_5$	$x + u_6 + w_1$	$u_6 + w_4$
$S_3$	$x + w_3$	$x + u_1 + w_1$	$u_3 + w_4$	$u_1 + w_7$	$x + u_3 + w_6$	$x + u_7 + w_2$	$u_7 + w_5$
$S_4$	$x + w_4$	$x + u_2 + w_2$	$u_4 + w_5$	$u_2 + w_1$	$x + u_4 + w_7$	$x + u_1 + w_3$	$u_1 + w_6$
$S_5$	$x + w_5$	$x + u_3 + w_3$	$u_5 + w_6$	$u_3 + w_2$	$x + u_5 + w_1$	$x + u_2 + w_4$	$u_2 + w_7$
$S_6$	$x + w_6$	$x + u_4 + w_4$	$u_6 + w_7$	$u_4 + w_3$	$x + u_6 + w_2$	$x + u_3 + w_5$	$u_3 + w_1$
$S_7$	$x + w_7$	$x + u_5 + w_5$	$u_7 + w_1$	$u_5 + w_4$	$x + u_7 + w_3$	$x + u_4 + w_6$	$u_4 + w_2$

Table 4: The Subspaces  $S_i$  Formed from Equation (5)

So we have a partition  $\mathcal{P}^*$  of  $V_7(2)$  of type  $\langle 1, 16, 0, 0 \rangle$  that consists of  $\{0, x\} \oplus U$ ,  $A_3$ ,  $A_4$ , and the 14 subspaces produced by equations (5) and (6).

Note that the set of entries of  $W^j$  equals  $W$ , and the set of entries of  $A_i^j$  equals  $A_i$ . So from each of equations (5) and (6), we can also obtain two sets of 7 subspaces of dimension 2 and 1 of the 3-dimensional subspaces  $A_i$  in a similar manner. These sets of 2-dimensional subspaces are described by the following equations.

$$(x + A_0^6) + A_1^1 = (x + A_2^5) \quad \text{and} \quad (x + W^3) + (x + A_3^1) = A_5^1 \quad (7)$$

$$A_0^2 + (x + A_1^6) = (x + A_6^7) \quad \text{and} \quad W^1 + (x + A_4^1) = (x + A_5^1) \quad (8)$$

In Examples 3–7 below, we use this partition  $\mathcal{P}^*$  to construct all the desired partitions in categories 5–20 of Table 3. The relevant example number is indicated in Table 3.

**Example 3** There is a partition of type  $\langle 1, 10, 14, 0 \rangle$  (category 10) that consists of  $\{0, x\} \oplus U$ ,  $A_3$ ,  $A_4$ ,  $A_6$ , the 7 subspaces of dimension 3 produced by equation (6), and the 14 subspaces of dimension 2 produced by equation (7). In addition, there is a partition of type  $\langle 1, 4, 28, 0 \rangle$  (category 16) that consists of  $\{0, x\} \oplus U$ ,  $A_2$ ,  $A_3$ ,  $A_4$ ,  $A_6$  and the 28 subspaces of dimension 2 produced by equations (7) and (8). Moreover, because any three of the subspaces  $A_i$  are contained in the 6-dimensional space  $U \oplus W$ , they can be reconfigured into 7 subspaces of dimension 2. Thus these partitions yield partitions of types  $\langle 1, 7, 21, 0 \rangle$  (category 13) and  $\langle 1, 1, 35, 0 \rangle$  (category 19). Therefore all of the partitions in categories 4, 10, 13, 16, and 19 of Table 3 can be realized.  $\square$

**Example 4** Proposition 6.2 and Corollary 6.3 show that partitions of types  $\langle 1, 13, 7, 0 \rangle$  and  $\langle 1, 13, 6, 3 \rangle$  do not exist. Thus, in category 7, we must construct a partition of type  $\langle 1, 13, 5, 6 \rangle$ . Such a partition can be obtained from  $\mathcal{P}^*$  by reconfiguring the 3-dimensional subspaces  $S_1$ ,  $S_2$ , and  $S_3$  into 5 subspaces of dimension 2 and 6 subspaces of dimension 1. The nonzero vectors in the subspaces of dimension 2 are:

$$\begin{aligned} \{x + u_6 + w_6, u_6 + w_4, x + w_3\}, & & \{x + u_5 + w_7, x + u_2 + w_5, u_3 + w_4\}, \\ \{x + w_1, u_1 + w_2, x + u_1 + w_4\}, & & \{x + w_2, x + u_7 + w_7, u_7 + w_6\}, \end{aligned}$$

and

$$\{x + u_1 + w_1, x + u_3 + w_6, u_7 + w_5\}.$$

Applying this reconfiguration to  $\mathcal{P}^*$ , we obtain a partition of type  $\langle 1, 13, 5, 6 \rangle$  (category 7).  $\square$

**Example 5** A partition of type  $\langle 1, 15, 1, 4 \rangle$  (category 5) can be obtained from  $\mathcal{P}^*$  by reconfiguring a subspace of dimension 3 into 1 subspace of dimension 2 and 4 subspaces of dimension 1. The same method can be used to construct a partition in categories 11, 14, 17, and 20 with  $j = 0$  from the corresponding partition in the preceding category.

Because there is no partition of type  $\langle 1, 14, 3, 5 \rangle$  (Proposition 6.4), we can use the same technique to produce the necessary partition in category 6. If we reconfigure each of two 3-dimensional subspaces in  $\mathcal{P}^*$  into a subspace of dimension 2 and 4 subspaces of dimension 1, we obtain a partition of type  $\langle 1, 14, 2, 8 \rangle$  (category 6).

This technique cannot be used to produce a partition of type  $\langle 1, 12, 8, 4 \rangle$  (category 8), however, because there is no partition of type  $\langle 1, 13, 7, 0 \rangle$ . Instead, we reconfigure the 3-dimensional subspaces  $S_1$ ,  $S_2$ ,  $S_3$ , and  $S_4$  in  $\mathcal{P}^*$  into 8 subspaces of dimension 2 and 4 subspaces of dimension 1. The nonzero vectors in the subspaces of dimension 2 are:

$$\begin{aligned} \{x + w_1, u_1 + w_7, x + u_1 + w_3\}, & & \{u_6 + w_5, u_7 + w_6, u_2 + w_1\}, \\ \{x + u_7 + w_7, u_7 + w_5, x + w_4\}, & & \{x + u_6 + w_1, x + u_3 + w_6, u_4 + w_5\}, \\ \{x + u_6 + w_6, u_5 + w_3, x + u_1 + w_4\}, & & \{u_2 + w_3, x + u_2 + w_5, x + w_2\}, \\ \{x + u_1 + w_1, u_3 + w_4, x + u_7 + w_2\}, & \text{and} & \{x + u_2 + w_2, x + u_4 + w_7, u_1 + w_6\}. \end{aligned}$$

Thus we obtain a partition of type  $\langle 1, 12, 8, 4 \rangle$  (category 8).  $\square$

**Example 6** It is possible to reconfigure the subspaces  $S_1$ ,  $S_2$ ,  $S_3$ ,  $S_5$ , and  $S_6$  into 10 subspaces of dimension 2 and 5 subspaces of dimension 1. The nonzero vectors in the subspaces of dimension 2 are:

$$\begin{array}{ll}
\{u_1 + w_2, u_6 + w_5, u_5 + w_3\}, & \{x + u_7 + w_7, u_4 + w_3, x + u_5 + w_1\}, \\
\{x + w_3, x + u_1 + w_1, u_1 + w_7\}, & \{x + w_6, x + u_3 + w_5, u_3 + w_1\}, \\
\{u_2 + w_3, u_3 + w_4, u_5 + w_6\}, & \{x + w_5, x + u_3 + w_3, u_3 + w_2\}, \\
\{x + u_4 + w_4, x + u_5 + w_7, u_7 + w_5\}, & \{x + w_2, x + u_6 + w_6, u_6 + w_7\}, \\
\{u_7 + w_6, x + u_2 + w_5, x + u_6 + w_1\}, & \text{and } \{x + w_1, x + u_6 + w_2, u_6 + w_4\}.
\end{array}$$

Applying this reconfiguration to  $\mathcal{P}^*$ , we obtain a partition of type  $\langle 1, 11, 10, 5 \rangle$  (category 9).

In this partition, we can use equation (8) to reconfigure 7 subspaces of dimension 3 into 1 subspace of dimension 3 (which is  $A_2$ ) and 14 subspaces of dimension 2. Applying this to the partition of type  $\langle 1, 11, 10, 5 \rangle$  described in the preceding paragraph produces a partition of type  $\langle 1, 5, 24, 5 \rangle$  (category 15). The subspaces of dimension 3 in this partition include  $A_2$ ,  $A_3$ , and  $A_4$ . As in Example 3, they can be reconfigured into 7 subspaces of dimension 2 to yield a partition of type  $\langle 1, 2, 31, 5 \rangle$  (category 18).  $\square$

**Example 7** The 7 subspaces from equation (5) and  $A_3$  can be reconfigured into 17 subspaces of dimension 2 and 5 subspaces of dimension 1. The nonzero vectors in the subspaces of dimension 2 are:

$$\begin{array}{ll}
\{u_4 + w_3, x + w_1, x + u_4 + w_7\}, & \{u_7 + w_6, x + w_2, x + u_7 + w_7\}, \\
\{u_2 + w_5, x + w_3, x + u_2 + w_2\}, & \{u_1 + w_2, x + w_4, x + u_1 + w_1\}, \\
\{u_3 + w_2, x + w_5, x + u_3 + w_3\}, & \{u_3 + w_1, x + w_6, x + u_3 + w_5\}, \\
\{u_6 + w_2, x + w_7, x + u_6 + w_6\}, & \{u_5 + w_1, x + u_4 + w_4, x + u_7 + w_2\}, \\
\{u_7 + w_1, x + u_4 + w_6, x + u_5 + w_5\}, & \{u_2 + w_3, u_3 + w_4, u_5 + w_6\}, \\
\{u_1 + w_6, u_2 + w_1, u_4 + w_5\}, & \{u_6 + w_7, x + u_1 + w_3, x + u_5 + w_1\}, \\
\{u_4 + w_7, u_7 + w_5, u_5 + w_4\}, & \{u_6 + w_5, x + u_1 + w_4, x + u_5 + w_7\}, \\
\{u_5 + w_3, x + u_2 + w_4, x + u_3 + w_6\}, & \{u_2 + w_7, x + u_7 + w_3, x + u_6 + w_1\},
\end{array}$$

and

$$\{u_1 + w_4, u_3 + w_6, u_7 + w_3\}.$$

Applying this reconfiguration to  $\mathcal{P}^*$  produces a partition of type  $\langle 1, 8, 17, 5 \rangle$  (category 12).  $\square$

Thus Examples 3–7 produce all the desired partitions in categories 5–20.

## 6.4 Partitions of $V_7(2)$ Containing No Subspace of Dimension Greater Than 3

The partitions of  $V_7(2)$  that do not contain a subspace of dimension 4 or more are listed in categories 21–38 of Table 3.

By Proposition 6.5, the only realizable partition type in category 21 is  $\langle 17, 0, 8 \rangle$ . A partition of this type can be obtained from  $\mathcal{P}^*$  by reconfiguring the subspace of dimension 4 into 1 subspace of dimension 3 and 8 subspaces of dimension 1.

With two exceptions, a partition type in category  $k$  ( $22 \leq k \leq 38$ ) can be obtained from the corresponding partition in category  $k - 18$  by reconfiguring the subspace of dimension 4 into 5 subspaces of dimension 2, and then reconfiguring 2-dimensional subspaces into subspaces of dimension 1 as necessary. For example, a partition of type  $\langle 16, 3, 6 \rangle$  (category 22) can be obtained from  $\mathcal{P}^*$  by reconfiguring the subspace of dimension 4 into 5 subspaces of dimension 2 and then reconfiguring 2 subspaces of dimension 2 into 6 subspaces of dimension 1.

The exceptions arise in categories 24 and 25 because, by Propositions 6.4 and 6.2, there are no partitions of types  $\langle 1, 14, 3, 5 \rangle$  (category 6) and  $\langle 1, 13, 7, 3 \rangle$  (category 7). So, to complete our analysis, we must construct partitions of types  $\langle 14, 8, 5 \rangle$  (category 24) and  $\langle 13, 12, 0 \rangle$  (category 25).

**Example 8** We construct a partition of type  $\langle 14, 8, 5 \rangle$  (category 24) by reconfiguring the subspaces  $\{0, x\} \oplus U$ ,  $S_1$ , and  $S_2$  into 8 subspaces of dimension 2 and 5 subspaces of dimension 1. The nonzero vectors in the subspaces of dimension 2 are:

$$\begin{aligned} & \{u_3, u_5 + w_3, u_2 + w_3\}, & & \{x + u_7 + w_7, x + u_2 + w_5, u_6 + w_4\}, \\ & \{x + u_1, u_1 + w_2, x + w_2\}, & & \{u_6 + w_5, x + u_1 + w_4, x + u_5 + w_7\}, \\ & \{u_6, x + w_1, x + u_6 + w_1\}, & & \{x + u_2, x + u_6 + w_6, u_7 + w_6\}, \\ & \{x + u_3, x + u_5, u_2\}, & \text{and} & \{x + u_4, x + u_7, u_5\}. \end{aligned} \quad \square$$

**Example 9** We construct a partition of type  $\langle 13, 12, 0 \rangle$  (category 25) by reconfiguring the subspaces  $\{0, x\} \oplus U$ ,  $S_1$ ,  $S_2$ ,  $S_3$ , and  $S_4$  into 1 subspace of dimension 3 and 12 subspaces of dimension 2. The nonzero vectors in the subspace of dimension 3 are:

$$\{u_4, u_3, x, u_6, x + u_4, x + u_3, x + u_6\},$$

and the nonzero vectors in the subspaces of dimension 2 are:

$$\begin{aligned} & \{u_1, x + u_1 + w_4, x + w_4\}, & & \{u_1 + w_2, u_5 + w_3, u_6 + w_5\}, \\ & \{u_1 + w_6, x + u_5, x + u_6 + w_6\}, & & \{u_1 + w_7, x + u_1 + w_3, x + w_1\}, \\ & \{u_2, x + u_2 + w_2, x + w_2\}, & & \{u_2 + w_1, x + u_6 + w_1, x + u_7\}, \\ & \{u_2 + w_3, x + u_2, x + w_3\}, & & \{u_4 + w_5, u_5, u_7 + w_5\}, \\ & \{u_3 + w_4, x + u_1 + w_1, x + u_7 + w_2\}, & & \{u_7, x + u_4 + w_7, x + u_5 + w_7\}, \\ & \{u_7 + w_6, x + u_1, x + u_3 + w_6\}, & \text{and} & \{u_6 + w_4, x + u_2 + w_5, x + u_7 + w_7\}. \end{aligned} \quad \square$$

Thus all the partition types in categories 21–38 can be realized except the partition of type  $\langle 17, 1, 5 \rangle$ .

The results in Sections 3, 5, and 6 establish our main theorem.

## 7 Equivalence of Partitions

In this paper, we have concentrated on the type of a partition. Let us call partitions of  $V_n(2)$  of the same type,  $\mathcal{P} = \{V_1, V_2, \dots, V_k\}$  and  $\mathcal{P}'$ , *equivalent* if there is a change of basis  $T: V_n(2) \rightarrow V_n(2)$  such that  $\mathcal{P}' = \{T(V_1), T(V_2), \dots, T(V_k)\}$ . In conclusion, we note that partitions of the same type need not be equivalent. Using the notation of Section 5, we define partitions  $\mathcal{P}$  and  $\mathcal{P}'$  of  $V_6(2)$  as follows. The nonzero vectors in the subspaces of  $\mathcal{P}$  are

$$\begin{aligned} V_1 &= \{u_1, w_3, u_1 + w_3\}, & V_2 &= \{u_2, u_3, u_5\}, & V_3 &= \{u_4, u_7 + w_3, u_5 + w_3\}, \\ V_4 &= \{u_6, u_3 + w_3, u_4 + w_3\}, & V_5 &= \{u_7, u_2 + w_3, u_6 + w_3\}, & & \{w_1, w_2, w_4\}, \\ & \{w_5, u_6 + w_6, u_6 + w_1\}, & & \{w_6, u_3 + w_7, u_3 + w_2\}, & & \{w_7, u_1 + w_4, u_1 + w_5\}, \\ & \{u_1 + w_1, u_2 + w_4, u_4 + w_2\}, & & \{u_2 + w_2, u_5 + w_7, u_3 + w_6\}, & & \{u_4 + w_4, u_2 + w_5, u_1 + w_7\}, \\ & \{u_5 + w_5, u_7 + w_1, u_4 + w_6\}, & & \{u_7 + w_7, u_6 + w_2, u_2 + w_6\}, & & \{u_1 + w_2, u_5 + w_6, u_6 + w_7\}, \\ & \{u_3 + w_4, u_5 + w_2, u_2 + w_1\}, & & \{u_4 + w_5, u_5 + w_1, u_7 + w_6\}, & & \{u_7 + w_2, u_4 + w_1, u_5 + w_4\}, \\ & \{u_3 + w_5, u_4 + w_7, u_6 + w_4\}, & & \{u_7 + w_4, u_2 + w_7, u_6 + w_5\}, & & \{u_1 + w_6, u_3 + w_1, u_7 + w_5\}, \end{aligned}$$

and the nonzero vectors in the subspaces of  $\mathcal{P}'$  are

$$\begin{array}{lll}
\{u_1, w_1, u_1 + w_1\}, & \{u_2, w_2, u_2 + w_2\}, & \{u_3, w_3, u_3 + w_3\}, \\
\{u_4, w_4, u_4 + w_4\}, & \{u_5, w_5, u_5 + w_5\}, & \{u_6, w_6, u_6 + w_6\}, \\
\{u_7, w_7, u_7 + w_7\}, & \{u_1 + w_2, u_4 + w_6, u_2 + w_7\}, & \{u_2 + w_3, u_6 + w_7, u_7 + w_1\}, \\
\{u_3 + w_4, u_2 + w_6, u_5 + w_3\}, & \{u_4 + w_5, u_7 + w_3, u_5 + w_2\}, & \{u_5 + w_6, u_1 + w_4, u_6 + w_3\}, \\
\{u_1 + w_3, u_5 + w_7, u_6 + w_1\}, & \{u_2 + w_4, u_4 + w_7, u_1 + w_5\}, & \{u_3 + w_5, u_6 + w_2, u_4 + w_3\}, \\
\{u_7 + w_2, u_6 + w_4, u_2 + w_1\}, & \{u_2 + w_5, u_3 + w_7, u_5 + w_4\}, & \{u_3 + w_6, u_4 + w_1, u_6 + w_5\}, \\
\{u_5 + w_1, u_7 + w_4, u_4 + w_2\}, & \{u_1 + w_6, u_3 + w_1, u_7 + w_5\}, & \{u_1 + w_7, u_3 + w_2, u_7 + w_6\}.
\end{array}$$

Both  $\mathcal{P}$  and  $\mathcal{P}'$  are partitions of type  $\langle 21, 0 \rangle$ . It is easily checked that the subspaces  $V_1, V_2, \dots, V_5$  in  $\mathcal{P}$  can be reconfigured into a 4-dimensional subspace. A more tedious calculation shows that the span of no 2 subspaces in  $\mathcal{P}'$  contains 5 subspaces in  $\mathcal{P}'$ , and so no 5 subspaces in  $\mathcal{P}'$  can be reconfigured into a 4-dimensional subspace. Thus  $\mathcal{P}$  and  $\mathcal{P}'$  are not equivalent.

## References

- [1] A. Beutelspacher, Partitions of finite vector spaces: an application of the Frobenius number in geometry, *Arch. Math.* **31** (1978), 202–208.
- [2] A. Blinco, S. El-Zanati, G. Seelinger, P. Sissokho, L. Spence, and C. Vanden Eynden, On vector space partitions and uniformly resolvable designs, in preparation.
- [3] W. Clark and L. Dunning, Partial partitions of vector spaces arising from the construction of byte error control codes, *Ars Combin.* **33** (1992), 161–177.
- [4] S. El-Zanati, G. Seelinger, P. Sissokho, L. Spence, and C. Vanden Eynden, Partitions of finite vector spaces into subspaces, in preparation.
- [5] T. Bu, Partitions of a vector space, *Discrete Math.* **31** (1980), 79–83.
- [6] O. Heden, On partitions of finite vector spaces of small dimensions, *Arch. Math.* **43** (1984), 507–509.